

Homework 11

¶ 1. Prove the following:

- A space X is contractible if and only if X is homotopically equivalent to a singleton.
- Every contractible space is path-connected.
- Every retract in a contractible space is contractible. (Recall that $A \subset X$ is called a retract of X if there is a continuous map $r : X \rightarrow A$ such that $r(a) = a$ for every a in A .)

¶ 2. (a) Prove that if $A \subset X$ is a deformation retract, then A and X are homotopically equivalent.

- Prove that any non-empty, compact, convex subset of \mathbf{R}^m is a deformation retract of \mathbf{R}^m .

¶ 3. A mapping $f : X \rightarrow Y$ is null-homotopic if it is homotopic to a constant mapping.

For a space X , let $CX = X \times [0, 1]/X \times \{1\}$, the cone on X , obtained from the product $X \times [0, 1]$ by identifying all points in the subset $X \times \{1\}$.

- Prove that a continuous mapping $f : X \rightarrow Y$ is null-homotopic if and only if f admits a continuous extension to CX .
- Prove that two null-homotopic mappings $f, g : X \rightarrow Y$ need not be homotopic to each other.

¶ 4. Prove or give counterexamples:

- If $f, g : X \rightarrow Y$ are homotopic and $A \subset X$, then the restrictions $f|_A$ and $g|_A$ are homotopic as maps $A \rightarrow Y$.
- If $f, g : X \rightarrow Y$ are homotopic and $B \subset Y$ is such that $f(X) \subset B$ and $g(X) \subset B$, then the f and g are homotopic as mappings from X into B .
- Two mappings f, g of a space X into a product space $Y = Y_1 \times \cdots \times Y_n$ are homotopic if and only if the compositions $\pi_k \circ f$ and $\pi_k \circ g$ are homotopic for all $k = 1, \dots, n$.
- Two mappings f, g of a space X into a product space $Y = \prod_{\alpha \in A} Y_\alpha$ are homotopic if and only if the compositions $\pi_\alpha \circ f$ and $\pi_\alpha \circ g$ are homotopic for all $\alpha \in A$.

¶ 5. Let $f_1, g_1 : X \rightarrow Y$ and $f_2, g_2 : Y \rightarrow Z$. Suppose that there are homotopies $H_1 : f_1 \simeq g_1$ and $H_2 : f_2 \simeq g_2$, and let $H : X \times I \rightarrow Z$ be given by $H(x, t) = H_2(H_1(x, t), t)$. True or False: H is a homotopy from $f_2 \circ f_1$ to $g_2 \circ g_1$.

¶ 6. Let (X, A) and (Y, B) be two pairs of spaces. A mapping $f : (X, A) \rightarrow (Y, B)$ is a continuous mapping $f : X \rightarrow Y$ such that $f(A) \subset B$.

Two mappings $f, g : (X, A) \rightarrow (Y, B)$ are homotopic relative to A if there is a continuous mapping $H : X \times [0, 1] \rightarrow Y$ such that

$$\begin{aligned} H(x, 0) &= f(x) && \text{for all } x \text{ in } X \\ H(x, 1) &= g(x) && \text{for all } x \text{ in } X \\ H(a, t) &= f(a) = g(a) && \text{for all } a \text{ in } A \text{ and all } t \text{ in } [0, 1] \end{aligned}$$

Prove that “being homotopic relative to A ” is an equivalence relation on the set of all mappings $f : (X, A) \rightarrow (Y, B)$.

¶ 7. Let X be the subset of the plane \mathbf{R}^2 consisting of the line segment from $(0, 0)$ to $(1, 0)$ on the x -axis, the line segment from $(0, 0)$ to $(0, 1)$ on the y -axis, and the line segments $\{(1/n, y) \mid 0 \leq y \leq 1\}$. Let $A \subset X$ be the subspace $A = \{(0, 1)\}$.

- Prove that X is contractible.
- Prove that the identity $f = \text{id}_X$ and the constant map $g = c_{(0,1)}$ are homotopic. (Here $c_{(0,1)}(x) = (0, 1)$ for all x .)
- Prove that f and g are not homotopic relative to A .

¶ 8. Let D be an open subset of \mathbf{R}^n , let α be a path in D from x to y , and set $\delta = \inf \{|\alpha(s) - w| \mid w \in \partial D, 0 \leq s \leq 1\}$. Show that if β is any path in D from x to y and $|\alpha(s) - \beta(s)| \leq \delta$ for all $0 \leq s \leq 1$, then α and β are homotopic relative endpoints.

¶ 9. A path α in \mathbf{R}^m is a polygonal path if there is a partition $0 = s_0 < s_1 < \cdots < s_n = 1$ of the interval $[0, 1]$ such that $\alpha(s) = \frac{s_i - s}{s_i - s_{i-1}} \alpha(s_{i-1}) + \frac{s - s_{i-1}}{s_i - s_{i-1}} \alpha(s_i)$ on $[s_{i-1}, s_i]$, for $i = 1, \dots, n$. Prove that any path in an open subset D of \mathbf{R}^m is homotopic in D relative to endpoints to a polygonal path.

¶ 10. Let (X, d) be a compact metric space and let $a, b \in X$. Let $\prod(X; a, b)$ be the set of paths in X from a to b , endowed with the metric $D(\alpha, \beta) = \sup \{d(\alpha(s), \beta(s)) \mid 0 \leq s \leq 1\}$. Prove that two paths α, β in $\prod(X; a, b)$ are homotopic relative to endpoints if they are in the same path connected component of the metric space $\prod(X; a, b)$, that is, if there is a path in $\prod(X; a, b)$ from a to b .