

Problem 1. (a) Define the concept “ f is a continuous function at the point p .”

(b) Let $f : \mathbf{R} \rightarrow \mathbf{R}$ be the function given by $f(x) = x$ if x is rational, and $f(x) = -x$ if x is irrational. Prove that f is continuous only at $p = 0$.

Problem 2 (IV.1). Discuss the continuity of the function $f : \mathbf{R} \rightarrow \mathbf{R}$ if

$$(a) f(x) = \begin{cases} 0, & x < 0 \\ x, & x \geq 0 \end{cases}$$

$$(b) f(x) = \begin{cases} x^2, & x \neq 0 \\ 1, & x = 0 \end{cases}$$

$$(c) f(x) = \sqrt{|x|}$$

Problem 3 (IV.10). Discuss the continuity of the function $f : \mathbf{R}^2 \rightarrow \mathbf{R}$ if

$$(a) f(x, y) = \begin{cases} \frac{1}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

$$(b) f(x, y) = \begin{cases} \frac{xy}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

$$(c) f(x, y) = \begin{cases} \frac{xy^2}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

Problem 4. For a non-empty subset S of \mathbf{R}^k , let d_S be the function given by $d_S(x) = \text{g.l.b.}\{|x - y| \mid y \in S\}$, where $|x - y|$ is the distance between the vectors x and y . Prove that d_S is continuous on \mathbf{R}^k .

Problem 5. (a) Give an example of a continuous function on a closed set $E \subset \mathbf{R}$ that has no maximum.

(b) Give an example of a continuous function on a bounded set $F \subset \mathbf{R}$ that has no maximum.

Problem 6 (IV.7). Let I be an interval in \mathbf{R} and let $a \in I$. If f is a function whose domain contains $I \setminus \{a\}$, define

$$\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a} f_+(x)$$

where f_+ is the function with domain $I \cap (a, \infty)$ given by $f_+(x) = f(x)$. Similarly, define

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a} f_-(x)$$

where f_- is the function with domain $I \cap (-\infty, a)$ given by $f_-(x) = f(x)$. Prove that $\lim_{x \rightarrow a} f(x)$ exists if and only if both

$\lim_{x \rightarrow a^+} f(x)$ and $\lim_{x \rightarrow a^-} f(x)$ exist and are equal.

Problem 7 (IV.8). Let f be a real valued function defined on (a, ∞) , where $a > 0$ is some positive real number. Let

$\lim_{x \rightarrow \infty^+} f(x)$ be given by

$$\lim_{x \rightarrow \infty^+} f(x) = \lim_{y \rightarrow 0} g(y),$$

where $g : (0, 1/a) \rightarrow \mathbf{R}$ is given by $g(y) = f(1/y)$, if this latter limit exists.

Prove that $\lim_{x \rightarrow \infty^+} f(x)$ exists if and only if for any $\varepsilon > 0$ there exists a number $N \geq a$ such that $|f(x) - f(y)| < \varepsilon$ if $x, y > N$.

Problem 8 (IV.4). Let I, J be open intervals in \mathbf{R} and let $f : I \rightarrow J$ be a function that is strictly increasing (that is, if $x < y$, then $f(x) < f(y)$) and surjective. Prove that f is continuous.

Problem 9. Suppose that f is continuous on an open set S . Prove that if p is in S , then there exists $r > 0$ such that $f(B(p, r))$ is a bounded set.

Problem 10. True or false:

- (a) f is continuous if and only if f takes convergent sequences to convergent sequences (that is, if p_1, p_2, p_3, \dots is a sequence in $\text{dom}(f)$ that converges to a point in $\text{dom}(f)$, then $f(p_1), f(p_2), f(p_3), \dots$ converges).
- (b) f is continuous if and only if f takes Cauchy sequences to Cauchy sequences (that is, if p_1, p_2, p_3, \dots is a Cauchy sequence in $\text{dom}(f)$, then $f(p_1), f(p_2), f(p_3), \dots$ is a Cauchy sequence).

Problem 11. (a) Consider a sequence of closed intervals $I_1 = [a_1, b_1], I_2 = [a_2, b_2], \dots$. Suppose that $a_n \leq a_{n+1}$ and $b_{n+1} \leq b_n$ for all n . Prove that there exists a point x that is in every I_n .

(b) Prove that if $\text{Length } I_n \rightarrow 0$, then the point x in (a) is unique.

(c) Show that this conclusion in Part (i) is false if we consider open intervals instead of closed intervals. Is it true if we consider open and bounded intervals?