

# Math 140

## Introductory Statistics

Next test on March 27

# At the Hockey games

Team	Arena Seating (thousands)
New Jersey Devils	19
New York Islanders	16
New York Rangers	18
Philadelphia Flyers	18
Pittsburgh Penguins	17

Construct the probability distribution for  $X$ , the probability for the total number of people that can attend two distinct games

# $P(X)$ , attendance at 2 hockey games

Teams	Maximum Attendance (thousands)
Devils/Islanders	$19 + 16 = 35$

So here we found  $X = 35$ ,

Find the rest and then calculate  
 $P(X)$  for all  $X$

How many possible  $X$  values will we have?

# P(X), attendance at 2 hockey games

Teams	Maximum Attendance (thousands)
Devils/Islanders	19 + 16 = 35

How many possible X values will we have?  
We are selecting 2 arenas out of 5

The number of X values is

$$\binom{5}{2} = \frac{5!}{3! 2!} = \frac{5 * 4 * 3 * 2 * 1}{3 * 2 * 1 * 2 * 1} = 10$$

# $P(X)$ , attendance at 2 hockey games

Teams	Maximum Attendance (thousands)
Devils/Islanders	$19 + 16 = 35$
Devils/Rangers	$19 + 18 = 37$
Devils/Flyers	$19 + 18 = 37$
Devils/Penguins	$19 + 17 = 36$
Islanders/Rangers	$16 + 18 = 34$
Islanders/Flyers	$16 + 18 = 34$
Islanders/Penguins	$16 + 17 = 33$
Rangers/Flyers	$18 + 18 = 36$
Rangers/Penguins	$18 + 17 = 35$
Flyers/Penguins	$18 + 17 = 35$

# $P(X)$ , attendance at 2 hockey games

Total Possible Attendance (thousands), $x$	Probability, $p$
33	0.1
34	0.2
35	0.3
36	0.2
37	0.2

What is the probability that attendance is at least 36,000?

# $P(X)$ , attendance at 2 hockey games

Total Possible Attendance (thousands), $x$	Probability, $p$
33	0.1
34	0.2
35	0.3
36	0.2
37	0.2

$P(X > \text{or} = 36) = 0.2 + 0.2 = 0.4$     40% probability

What do you think is the average number of cars per household?

Vehicles per Household, $x$	Proportion of Households, $p$
0	0.087
1	0.331
2	0.381
3	0.201



What do you think is the average number of cars per household?

Vehicles per Household, $x$	Proportion of Households, $p$
0	0.087
1	0.331
2	0.381
3	0.201

$$\text{Average} = 0 * 0.087 + 1 * 0.331 + 2 * 0.381 + 3 * 0.201 = 1.696$$

# In a more abstract way

Average = Sum over all  $(X * P(X))$

The mean of a probability distribution for the random variable  $X$  is called its **expected value** and is denoted by  $\mu_x$  or  $E(X)$ .

The mean gets called **expected value** of the distribution  
Sometimes indicated as  $E(X)$  or  $\mu_x$

# Getting insured

The probability that you get burglarized is  
26.9 per 1000 people

Outcome	Payout, $x$	Probability, $p$	$x \cdot p$
No burglary	0	0.9731	0
Burglary	5000	0.0269	134.50


What the company will pay is \$5,000

# Getting insured

The expected payout the company will pay is

0 if there are no burglaries  
\$5000 if there is a burglary

For each policy we will pay on average  
 $0 \cdot 0 + 5000 \cdot 0.0269$



Outcome	Payout, $x$	Probability, $p$	$x \cdot p$
No burglary	0	0.9731	0
Burglary	5000	0.0269	134.50

$$= \text{Sum} (X * P(X)) = 134.50$$

# Getting insured

The company will break even if they charge 134.50

97% of the time people will pay,  
not be burglarized and not get anything,

The rest of the time they will be burglarized and get 5K

The company breaks even, since

$$E(X) = X P(X)$$

What everyone pays = What the company pays

# The Wisconsin lottery

Winnings, $x$	Probability, $p$
\$1	1/10
\$2	1/14
\$3	1/24
\$18	1/200
\$50	1/389
\$150	1/20,000
\$900	1/120,000

A ticket costs \$1.  
Why don't the probabilities add to one?  
What is missing?

# The Wisconsin lottery

Winnings, $x$	Probability, $p$
\$1	1/10
\$2	1/14
\$3	1/24
\$18	1/200
\$50	1/389
\$150	1/20,000
\$900	1/120,000

What is the expected value for the probability distribution?

# The Wisconsin lottery

Winnings, $x$	Probability, $p$
\$1	1/10
\$2	1/14
\$3	1/24
\$18	1/200
\$50	1/389
\$150	1/20,000
\$900	1/120,000

$$P(0) = 1 - \text{Sum of the above} = 0.7793$$

$$\mu_x = 0.6014$$

If we spend \$1 we will get back, on average, 60 cents.



# The Wisconsin lottery

We are given 10 tickets.  
How much do we expect to win?

0.6014 for each ticket

$10 * 0.6014 = \$6.01$  for 10 tickets

# Calculate the expected value

Value, $x$	Frequency, $f$
5	12
6	23
8	15

Value, $x$	Proportion, $f/n$
5	0.24
6	?
8	?

There are  
 $n=50$   
outcomes

# Calculate the expected value

Value, $x$	Frequency, $f$
5	12
6	23
8	15

There are  
 $n=50$   
outcomes

Value, $x$	Proportion, $f/n$
5	0.24
6	?
8	?

$$\mu_x = \sum \left( x \frac{f}{n} \right)$$

$\sim 6.3$

# Homework

Page 286 E13, E11

## 6.2 Variances of Probability distributions

$$\sigma_n^2 = \sum \frac{(x - \bar{x})^2}{n}$$

$$SD = \sqrt{\sigma_n^2} = \sqrt{\sum \frac{(x - \bar{x})^2}{n}}$$

For a set of data - note the n instead of n-1

## 6.2 Variances of Probability distributions

For a probability distribution  $P(X)$

$$\sigma^2_x = \sum \frac{(x - \bar{x})^2}{n}$$

$$SD = \sqrt{\sigma^2_n} = \sqrt{\sum \frac{(x - \bar{x})^2}{n}}$$

# Try calculating the variance and SD

Winnings, $x$	Probability, $p$
\$1	1/10
\$2	1/14
\$3	1/24
\$18	1/200
\$50	1/389
\$150	1/20,000
\$900	1/120,000

Recall, the average is 60.14 cents or 0.6014 dollars

# Try calculating the variance and SD

Winnings, $x$	Probability, $p$
\$1	1/10
\$2	1/14
\$3	1/24
\$18	1/200
\$50	1/389
\$150	1/20,000
\$900	1/120,000

The way to think about this is: for each deviation from the mean, what is the probability? For example  
The contribution for the \$1 winning is

$$(1 - 0.6014)^2 * 1/10$$



# Try calculating the variance and SD

Winnings, $x$	Probability, $p$
\$1	1/10
\$2	1/14
\$3	1/24
\$18	1/200
\$50	1/389
\$150	1/20,000
\$900	1/120,000

So all together, the variance is

$$(1 - 0.6014)^2 * 1/10 + (2 - 0.6014)^2 * 1/14 + \text{etc etc} = 16.32$$

The standard deviation is its square root = \$4.04

# Variances

## Formula for the Variance of a Probability Distribution

The variance of a discrete probability distribution is given by

$$\text{Var}(X) = \sigma_X^2 = \sum (x - \mu_X)^2 \cdot p$$

Here,  $p$  is the probability that the random variable  $X$  takes on the specific value  $x$  and  $\mu_X$  is the expected value. To get the standard deviation, take the square root of the variance.

# Significance

The average payout is \$0.60  
With a SD of \$4.04

This means on average you win little  
but there is a chance of winning a lot  
(that is why the SD is large)

# Let's triple the lottery

Original Winnings, $x$	Winnings in Special Promotion, $3x$	Probability, $p$
\$0	\$0	0.7793
\$1	\$3	1/10
\$2	\$6	1/14
\$3	\$9	1/24
\$18	\$54	1/200
\$50	\$150	1/389
\$150	\$450	1/20,000
\$900	\$2700	1/120,000

Calculate mean and standard deviation  
Compare to original lottery

# Let's triple the lottery

Original lottery winnings \$0.6014

New winnings \$1.804

Is this smart on Wisconsin's part?

Original SD = \$4.04

New SD = \$12.12

How are they related?

They are both multiplied by 3!

$$SD = \sigma_X$$

# Let's triple the lottery

$\mu_x$  turns into  $3 \mu_x$

$\sigma_x$  turns into  $3 \sigma_x$

This is true in general.  
Multiplying data will rescale average and SD  
of the distribution

What if I had decided to add  
50 cents to each winning?

# Let's add 50 cents to the payout

$\mu_x$  turns into  $0.50 + \mu_x$

$\sigma_x$  stays  $\sigma_x$

Just like rescaling and recentering  
So, adding C and multiplying by D gives

$\mu_x$  turns into  $\mu_{C+DX} = C + D * \mu_x$

$\sigma_x$  turns into  $\sigma_{C+DX} = D * \sigma_x$

# In general

## Linear Transformation Rule: The Effect of a Linear Transformation of $X$ on $\mu_X$ and $\sigma_X$

Suppose you have a probability distribution for random variable  $X$ , with mean  $\mu_X$  and standard deviation  $\sigma_X$ . If you transform each value by multiplying it by  $d$  and then adding  $c$ , where  $c$  and  $d$  are constants, then the mean and the standard deviation of the transformed values are given by

$$\mu_{c+dX} = c + d\mu_X$$

$$\sigma_{c+dX} = |d| \sigma_X$$

Now, this was for TRIPLING the lottery

What if we kept the same lottery and bought 3 tickets?



# What do you think?

If every time I play my average payout is \$0.6014  
What do I get after buying 3 tickets?

# What do you think?

If every time I play my average payout is \$0.6014  
What do I get after buying 3 tickets?

$$\text{Duh} - 3 * 0.6014 = 1.804!$$

Just like before!

It does not matter if I triple the lottery or  
if I buy three tickets, the result is the same.

My take-home on average is tripled.

# What do you think?

We can conclude that when we select three items from the same distribution we find

$$\mu_{3,X} = 3 \mu_x = \mu_x + \mu_x + \mu_x$$

In general, for different distributions we get

$$\mu_{X,Y} = \mu_x + \mu_y$$

# Two tickets from two lotteries

Let's buy a ticket from the lottery of  
California and of Texas

California  $\mu_x = \$0.50$

Texas  $\mu_y = \$0.75$

What are the expected total winnings?

$$\mu_{CA,TX} = \mu_{CA} + \mu_{TX} = \$0.50 + \$0.75 = \$1.25$$

# What about the standard deviation?

We will look at that next time.

# Practice and hk

Page 286

E11, E12, E13, E14, E15, E16

Page 295

P10, P13, E19, E20,