



**The Oil and Gas Resource
Potential of the
Arctic National Wildlife Refuge
1002 Area, Alaska**

**ANWR Assessment Team
Open File Report 98-34
U.S. Geological Survey**




Table of Contents

Map of Petroleum Plays

Table of Contents

Instructions for Viewing and Printing

Fact Sheet (Executive Summary)

Assessment Overview

Assessment Results

Play Descriptions

Topset Play

Turbidite Play

Wedge Play

Thomson Play

Kemik Play

Undeformed Franklinian Play

Deformed Franklinian Play

Thin-Skinned Thrust-Belt Play

Ellesmerian Thrust-Belt Play

Niguanak-Aurora Play

Help K. Takahashi and P. Nelson

K. Bird and D. Houseknecht

AO K. Bird

RS J. Schuenemeyer

P1 D. Houseknecht and C. Schenk

P2 D. Houseknecht and C. Schenk

P3 D. Houseknecht and C. Schenk

P4 C. Schenk and D. Houseknecht

P5 C. Schenk and D. Houseknecht

P6 J. Kelley and others

P7 J. Grow and others

P8 W. Perry and others

P9 J. Grow and others

P10 J. Grow and others

Methodology

ME J. Schuenemeyer

Assessment Definitions

DF R. Charpentier

Economic Analysis

EA E. Attanasi

General Geology and Physical Setting

Geographic and Geologic Setting

GG K. Bird

Image Gallery

IG K. Takahashi

Field Studies, 1995-1997

FS C. Schenk and others

Stratigraphy and Age Dating

Formation Properties

FP P. Nelson and K. Bird

Basement Rocks

BR J. Kelley

Carboniferous and Older Carbonates

CC J. Dumoulin

Biostratigraphic Framework

BI W. Poag

Brookian Sequences

BS D. Houseknecht and C. Schenk

Thomson and Kemik Sandstones

TK C. Schenk and D. Houseknecht

Sequence Stratigraphy of the Pebble Shale Unit

SS J. Macquaker and others

Structural Setting

Structural Modeling

SM F. Cole and others

Balanced Cross Section

BC T. Moore

Brookian Deformation

BD C. Potter and others

Niguanak and Aurora Structures

NA J. Grow and others

Geophysics

Seismic Processing

SP M. Lee and others

Aeromagnetic Modeling

AM J. Phillips

Gravity Modeling

GR R. Saltus and others

Geochemistry

Oil Analyses

OA P. Lillis and others

Petroleum Occurrence and Timing of Migration

FI R. Burruss

Vitrinite Reflectance

VR K. Bird and others

Source Rocks

SR M. Keller and others

Basin Evolution

BE E. Rowan

Fluid Flow Modeling

FF D. Hayba and others

Petroleum Systems

PS L. Magoon and others

Hydrocarbon Potential of Brookian Strata

HG D. Houseknecht and D. Hayba

Production Analysis

PA J. Quinn

Properties of Water and Rocks

Streamwater Analyses

SA B. Wang

Well Data and Well Plots

WL P. Nelson

Petrophysical Properties

PP P. Nelson

Appendixes

Field Notes and Location Maps

Methodology - Computer Code

Oil Analyses – Tables and Chromatograms

Vitrinite Reflectance - Tables

Petroleum Systems – Tables

Streamwater Analyses - Tables

Well Data - Files

CM C. Molenaar

ME J. Schuenemeyer

OA P. Lillis and others

VR K. Bird and others

PS L. Magoon and others

SA B. Wang

WL P. Nelson

Arctic National Wildlife Refuge, 1002 Area, Petroleum Assessment, 1998

INTRODUCTION

The Alaska National Interest Lands Conservation Act (1980) established the Arctic National Wildlife Refuge (ANWR) (fig. 1) as a wildlife refuge. In section 1002 of that act, Congress deferred a decision regarding future management of the 1.5-million-acre coastal plain (“1002 area”) in recognition of the area’s potentially enormous oil and gas resources and its importance as wildlife habitat. A report on the resources (including petroleum) of the 1002 area was submitted in 1987 to Congress by the Department of the Interior (DOI). Since completion of that report, numerous wells have been drilled and oil fields discovered near ANWR (fig. 2), new geologic and geophysical data have become available, seismic processing and interpretation capabilities have improved, and the economics of North Slope oil development have changed significantly.

The U.S. Geological Survey (USGS) commonly is asked to provide the Federal Government with timely scientific information in support of decisions regarding land management, environmental quality, and economic and strategic policy. To do so, the USGS must anticipate issues most likely to be the focus of policymakers in the future. Anticipating the need for scientific information and considering the decade-old perspective of the petroleum resource estimates included in the 1987 Report to Congress, the USGS has reexamined the geology of the ANWR 1002 area and prepared a new petroleum resource assessment.




Figure 1. Map of northern Alaska and nearby parts of Canada showing locations of the Arctic National Wildlife Refuge (ANWR), the 1002 area, and the National Petroleum Reserve—Alaska (NPRA). Locations of known petroleum accumulations and the Trans-Alaska Pipeline System (TAPS) are shown, as well as summaries of known petroleum volumes in northern Alaska and the Mackenzie River delta of Canada. BBO, billion barrels of oil (includes cumulative production plus recoverable resources); TCFG, trillion cubic feet of gas recoverable resources.




Figure 2. Map of the ANWR 1002 area. Dashed line labeled Marsh Creek anticline marks approximate boundary between undeformed area (where rocks are generally horizontal) and deformed area (where rocks are folded and faulted). Boundary is defined by Marsh Creek anticline along western half of dashed line and by other geologic elements along eastern half of dashed line. Exploration wells are coded to show whether information from them was available for the 1987 USGS assessment of in-place petroleum resources. Dashed red line shows the offshore extent of the entire assessment area.

ASSESSMENT PROJECT

The new assessment involved 3 years of study by 40 USGS scientists, who coordinated work with colleagues in other Federal agencies, Alaska State agencies, and several universities. New field studies were conducted, new well and sample data were analyzed, and new geophysical data were acquired. Perhaps most importantly, all 1,400 miles of seismic data that had been collected by a petroleum-industry consortium in 1984 and 1985 were reprocessed and reinterpreted. Collection of seismic data within ANWR requires an act of Congress, and these are the only seismic data ever collected within the 1002 area. All this information was integrated as basic input to the petroleum assessment. The term “petroleum” is used herein to include crude oil, natural gas, and natural gas liquids. Although all petroleum commodities were assessed, results reported in this Fact Sheet are for crude oil only because it determines the economic viability of resources on the North Slope. Results for the other commodities are reported in a CD-ROM (USGS Open-File Report 98-34).

ASSESSMENT METHODOLOGY

The methodology used in this assessment is slightly modified from that used in the 1987 assessment of this area when the USGS estimated in-place petroleum resources for the DOI Report to Congress; the methodology is also slightly modified from that used in the USGS assessment of the National Petroleum Reserve—Alaska (NPR) (1979–1980). Ten petroleum *plays* were defined as the initial step of the assessment (fig. 3). A play is a volume of rock that contains similar geological parameters (such as petroleum charge, reservoir, and trap) that determine petroleum potential. In keeping with the USGS responsibility for assessing the petroleum potential of all onshore and State water areas of the United States, the total play area considered was extended to the 3-mile boundary between State and Federal jurisdiction. Thus, in addition to the Federal lands of the ANWR 1002 area, this assessment includes resources associated with State waters and Native lands (fig. 2).

For each play, distributions of the number and size of potential petroleum accumulations were estimated based on a probabilistic range of values for certain geological attributes, such as reservoir thickness and porosity. These distributions were restricted to potential accumulations larger than 50 million barrels of oil (MMBO) in-place so that the assessment would not be influenced by smaller accumulations that are non-economic in most cases on the North Slope.

The resulting distributions were subjected to a geologic risking procedure designed to weigh the likelihood that petroleum charge, reservoir, and trap conditions were sufficient to generate a 50-MMBO in-place accumulation. In turn, a probabilistic estimate of *in-place* petroleum resources was calculated based on the risked distributions of size and number of potential petroleum accumulations in each play. A recovery factor appropriate to each play was applied to the estimates of in-place petroleum resources to calculate *technically recoverable* petroleum resources (fig. 4).

Estimates for each play were aggregated to calculate total technically recoverable petroleum resources for the entire assessment area, the 1002 area, and the undeformed and deformed parts of the 1002 area (table 1). Costs associated with discovering and recovering petroleum resources, including the costs of constructing pipelines to transport the petroleum, were then applied to estimate *economically recoverable* petroleum resources (fig. 4).

The assessment methodology yields results that include probabilistic expressions of uncertainty, as illustrated schematically in figure 4. To stress the importance of this uncertainty, results reported here include 95- and 5-percent probabilities, in addition to mean value. The 95- and 5-percent probabilities are considered reasonable minimum and maximum values, and the mean is the average or expected value.

The 95-percent probability level refers to 19 in 20 chances; the 5-percent probability level refers to 1 chance in 20 that the amounts shown will be at least that large.





Figure 3. Summary of ages, names, and rock types present in the ANWR 1002 area. The occurrence of recoverable petroleum in these rock formations outside the ANWR 1002 area is indicated by green and red circles. Gray bars at right indicate the 10 petroleum plays assessed in this study and their corresponding rock formations (to the left). Note grouping of plays according to deformed and undeformed areas as shown in figure 2.

Charge.—Occurrence of conditions of petroleum generation and migration adequate to cause an accumulation of the minimum size (50 million barrels of oil (MMBO) in-place).

Reservoir.—Occurrence of reservoir rocks of sufficient quantity and quality to permit containment of petroleum in volumes sufficient for an accumulation of the minimum size (50 MMBO in-place).

Trap.—Occurrence of those structures, pinch-outs, permeability changes, and similar features necessary for the entrapment and sealing of petroleum in an accumulation of the minimum size (50 MMBO in-place).

ASSESSMENT RESULTS

The total quantity of technically recoverable oil within the entire assessment area is estimated to be between 5.7 and 16.0 billion barrels (95-percent and 5-percent probability range), with a mean value of 10.3 billion barrels. Technically recoverable oil within the ANWR 1002 area (excluding State and Native areas) is estimated to be between 4.3 and 11.8 billion barrels (95- and 5-percent probability range), with a mean value of 7.7 billion barrels (table 1).

Quantities of technically recoverable oil are not expected to be uniformly distributed throughout the ANWR 1002 area. The undeformed area (fig. 2) is estimated to contain between 3.4 and 10.2 billion barrels of oil (BBO) (95- and 5-percent probability), with a mean of 6.4 BBO. The deformed area (fig. 2) is estimated to contain between 0 and 3.2 BBO (95- and 5-percent probability), with a mean of 1.2 BBO.

Figure 5 shows the expected numbers of accumulations and volumes of technically recoverable oil relative to sizes of accumulations. Economic viability of an individual accumulation depends on the price of oil, availability of recovery technology, and the proximity of infrastructure (pipelines, etc.).

An important threshold for economic viability is an accumulation size that is sufficiently large to warrant development, and, in the past, a minimum economic field size for the study area has been considered to be about 400 MMBO recoverable. In recent years, however, North Slope infrastructure has expanded and industry has applied innovative technologies to dramatically reduce costs. As a result, development of fields as small as 150 MMBO recoverable is occurring, even at relatively low well-head prices.

If one considers only the western part of the ANWR 1002 area—plays of the undeformed area (fig. 2)—and assumes, rather conservatively, a minimum commercially developable field size of 512 million barrels recoverable, then about 2.6 BBO is expected to be economically recoverable from about three accumulations (fig. 5; sum of resources represented by all accumulations larger than 512 MMBO). Furthermore, this part of the assessment area is relatively confined geographically, so that much smaller accumulations (as small as 100 MMBO) may be commercial if they can be developed jointly with other accumulations or as satellite fields. For the deformed area (fig. 2), considerable uncertainty exists about using a similar 512-MMBO minimum field size because this area is much farther from infrastructure. If a similar minimum field size is assumed, one would expect about 600 MMBO economically recoverable from one field (fig. 5).

In-place resources.—The amount of petroleum contained in accumulations of at least 50 MMBO without regard to recoverability.

Technically recoverable resources.—Volume of petroleum representing that proportion of assessed in-place resources that may be recoverable using current recovery technology without regard to cost.

Economically recoverable resources.—That part of the technically recoverable resource for which the costs of discovery, development, and production, including a return to capital, can be recovered at a given well-head price.

Table 1. Estimates of volumes of technically recoverable oil in various parts of the ANWR assessment study area.

[ANWR, Arctic National Wildlife Refuge. All reported oil volumes in millions of barrels. Basic statistical principles determine that mean values can be added and subtracted but F_{95} and F_{05} values cannot (e.g., means for the undeformed and deformed parts of the ANWR 1002 area sum to the mean for the total ANWR 1002 area, but F_{95} and F_{05} values do not). F_{95} , 95-percent probability level; F_{05} , 5-percent probability level]

Part of study area	Volume of oil, in millions of barrels		
	F_{95}	Mean	F_{05}
Entire assessment area ¹	5,724	10,322	15,955
ANWR 1002 area (Federal), TOTAL	4,254	7,668	11,799
Undeformed part	3,403	6,420	10,224
Deformed part	0	1,248	3,185

¹ Includes 1002 area shown on figure 2, Native lands, and adjacent State water areas within 3-mile boundary (see fig. 2).




Figure 4. Schematic graph illustrating petroleum volumes and probabilities. Curves represent categories of oil in assessment. An example of how one reads this graph is illustrated by the blue and orange lines projected to the curve for economically recoverable oil. There is a 95-percent chance (i.e., probability, F_{95}) of at least volume V_1 of economically recoverable oil, and there is a 5-percent chance (F_{05}) of at least volume V_2 of economically recoverable oil.



Typical view of the ANWR 1002 area coastal plain.




Figure 5. A, Histogram showing the expected (mean) number of petroleum accumulations estimated to exist in various size categories of technically recoverable oil resources in this assessment. For each size category, histogram bar on left is for undeformed area and bar on right is for deformed area. Each histogram bar is divided into Federal (1002 area) and non-Federal portions. The histogram is read as follows: It is estimated that the undeformed area contains approximately three accumulations containing between 512 and 1,024 million barrels of technically recoverable oil; two of those accumulations are under Federal jurisdiction and one is non-Federal. Adding the accumulations in the undeformed area to those of the deformed area for each size category gives the total number of those sized accumulations for the entire assessment area (1002 area plus non-1002).

B, Histogram showing the expected (mean) volume of oil estimated to exist in each accumulation-size category of technically recoverable oil resources. For each size category, histogram bar on left is for undeformed area and bar on right is for deformed area. Each histogram bar is divided into Federal and non-Federal portions. The histogram is read as follows: It is estimated that the undeformed area contains approximately 1,800 million barrels of technically recoverable oil in accumulations containing between 512 and 1,024 million barrels of technically recoverable oil; approximately 1,300 MMBO is Federal and approximately 500 MMBO is non-Federal. Adding the volume of oil in the undeformed area to that of the deformed area for each size category gives the total volume of those sized accumulations for the entire assessment area (1002 area plus non-1002).

TECHNICALLY RECOVERABLE OIL GROUPED BY ACCUMULATION-SIZE CLASS, IN MILLIONS OF BARRELS



Photograph of oil-stained sandstone near crest of Marsh Creek anticline, 1002 area.

COMPARISON WITH PREVIOUS ASSESSMENTS

One cannot make a meaningful comparison with previous assessments without knowledge of differences in assessment methodology, assumptions, and data. That information is not always available for the previous assessments of the ANWR 1002 area. Among previous assessments of ANWR 1002 area petroleum resources, only the 1987 USGS assessment of in-place resources is directly comparable. The technically and economically recoverable petroleum resource estimates cannot be compared directly because different methods were used in preparing those parts of the 1987 Report to Congress. The current assessment shows an overall increase in estimated in-place oil resource when compared to the 1987 assessment. Ranges are 11.6 to 31.5 BBO versus 4.8 to 29.4 BBO, (95- and 5-percent probabilities) and mean values are 20.7 BBO versus 13.8 BBO (current assessment compared to 1987 assessment). The increase results in large part from improved resolution of reprocessed seismic data, which allowed the identification of many more potential petroleum accumulations in parts of the area, and analog information provided by recent nearby oil discoveries.

Another significant change is in the geographic distribution of resources. In the 1987 assessment, about 75 percent of the mean estimated in-place oil was in the southeastern or deformed area and only 25 percent was in the northwestern or undeformed area (fig. 2). In the current assessment, nearly 85 percent of the in-place oil is in the undeformed area and only about 15 percent is within the deformed area. Again, the reason is largely related to improved resolution of the seismic data, especially in the undeformed area where, in various plays, it allowed the identification of many more potential petroleum accumulations than were previously thought to exist. The southeastern area—with only a single well offshore and complex geology onshore—carries great uncertainty. Further, part of that area considered oil prospective in 1987 is now considered prospective only for gas because of new understanding of the thermal history of the rocks.

SUMMARY

In anticipation of the need for scientific support for policy decisions and in light of the decade-old perspective of a previous assessment, the USGS has completed a reassessment of the petroleum potential of the ANWR 1002 area. This was a comprehensive study by a team of USGS scientists in collaboration on technical issues (but not the assessment) with colleagues in other agencies and universities. The study incorporated all available public data and included new field and analytic work as well as the reevaluation of all previous work.

Using a methodology similar to that used in previous USGS assessments in the ANWR and the NPRA, this study estimates that the total quantity of technically recoverable oil in the 1002 area is 7.7 BBO (mean value), which is distributed among 10 plays. Using a conservative estimate of 512 million barrels as a minimum commercially developable field size, then about 2.6 BBO of oil distributed in about three fields is expected to be economically recoverable in the undeformed part of the 1002 area. Using a similar estimated minimum field size, which may not be conservative considering the increased distance from infrastructure, the deformed area would be expected to have about 600 MMBO in one field.

The amounts of in-place oil estimated for the 1002 area are larger than previous USGS estimates. The increase results in large part from improved resolution of reprocessed seismic data and geologic analogs provided by recent nearby oil discoveries.

The U.S. Geological Survey is solely responsible for the input and results of this assessment. The USGS acknowledges the cooperation of the U.S. Fish and Wildlife Service, Bureau of Land Management, Minerals Management Service, and the Alaska Department of Natural Resources (Geological and Geophysical Surveys, Division of Oil and Gas, and Oil and Gas Conservation Commission), which provided access to data as well as feedback on geology and methodology.

For further information and to request a CD-ROM (USGS Open-File Report 98-34) containing detailed results and supporting scientific documentation, send e-mail to:

gd-anwr@usgs.gov

or contact:

Kenneth J. Bird kbird@usgs.gov (650) 329-4907

David W. Houseknecht dhouse@usgs.gov (703) 648-6470

