

## 4.4 Matrices: Basic Operations

- Addition and subtraction of matrices
- Product of a number  $k$  and a matrix  $M$
- Matrix Product.

### Addition and Subtraction of matrices

- To add or subtract matrices, they must be of the same size  $m \times n$ .
- To add matrices of the same size, add their corresponding entries.

$$A + B = \begin{bmatrix} a_{ij} + b_{ij} \end{bmatrix} \quad \begin{pmatrix} 1 & 2 \\ 5 & 2 \end{pmatrix} + \begin{pmatrix} 4 & 5 \\ 9 & 1 \end{pmatrix} =$$

- To subtract matrices of the same order, subtract their corresponding entries. The general rule is as follows using mathematical notation:

$$A - B = \begin{bmatrix} a_{ij} - b_{ij} \end{bmatrix} \quad \begin{pmatrix} 1 & 2 \\ 5 & 2 \end{pmatrix} - \begin{pmatrix} 4 & 5 \\ 9 & 1 \end{pmatrix} =$$

## More examples:

$$\begin{bmatrix} 4 & -3 & 1 \\ 0 & 5 & -2 \\ 5 & -6 & 0 \end{bmatrix} + \begin{bmatrix} -1 & 2 & 3 \\ 6 & -7 & 9 \\ 0 & -4 & 8 \end{bmatrix}$$

$$\begin{bmatrix} 4 & -3 & 1 \\ 0 & 5 & -2 \\ 5 & -6 & 0 \end{bmatrix} - \begin{bmatrix} -1 & 2 & 3 \\ 6 & -7 & 9 \\ 0 & -4 & 8 \end{bmatrix}$$

$$\begin{bmatrix} 4 & -3 & 1 \\ 0 & 5 & -2 \\ 5 & -6 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 5 \\ 3 & 7 \\ 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 6 & 3 \end{bmatrix} - \begin{bmatrix} -2 & 3 & 1 \end{bmatrix}$$

## Scalar Multiplication

- The **scalar product** of a number  $k$  and a matrix  $\mathbf{A}$  is the matrix denoted by  $k\mathbf{A}$ , obtained by multiplying each entry of  $\mathbf{A}$  by the number  $k$ .
- The number  $k$  is called a **scalar**.

$$k\mathbf{A} = \begin{bmatrix} ka_{ij} \end{bmatrix}$$

- **Example:**

$$(-1) \begin{bmatrix} -1 & 2 & 3 \\ 6 & -7 & 9 \\ 0 & -4 & 8 \end{bmatrix}$$

## More Examples of scalar multiplication & addition:

$$\begin{bmatrix} -1 & 2 & 3 \\ 6 & -7 & 9 \\ 0 & -4 & 8 \end{bmatrix} + 3 \begin{bmatrix} 1 & 4 & 2 \\ 0 & 4 & 1 \\ 2 & 5 & -1 \end{bmatrix} =$$

$$(2) \begin{bmatrix} -1 & 2 & 3 \\ 6 & -7 & 9 \\ 0 & -4 & 8 \end{bmatrix} + (0) \begin{bmatrix} 1 & 4 & 2 \\ 0 & 4 & 1 \\ 2 & 5 & -1 \end{bmatrix} =$$

## Alternate definition of subtraction of matrices:

- The definition of subtract of two real numbers  $a$  and  $b$  is

$$a - b = a + (-1)b$$

i.e.  $a$  plus negative  $b$ .

- We can define subtraction of matrices similarly:


If  $A$  and  $B$  are two matrices of the same dimensions, then

$$A - B = A + (-1)B,$$

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} - \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} + (-1) \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix} =$$

## Matrix product

- The method of multiplication of matrices is not as intuitive and may seem strange, although this method is extremely useful in many mathematical applications.
- Matrix multiplication was introduced by an English mathematician named Arthur Cayley (1821-1895) .
- We will see shortly how matrix multiplication can be used to solve systems of linear equations.



## Row by column multiplication

- 1X4 row matrix multiplied by a 4X1 column matrix: Notice the manner in which corresponding entries of each matrix are multiplied:

$$(1 \ 2 \ 3 \ 4) \begin{pmatrix} 5 \\ 6 \\ 7 \\ 8 \end{pmatrix} = 1 \cdot 5 + 2 \cdot 6 + 3 \cdot 7 + 4 \cdot 8 = 70$$

$$(4 \ 2 \ 3 \ 1) \begin{pmatrix} 6 \\ 3 \\ 5 \\ 2 \end{pmatrix} =$$

## Row by column multiplication

$1 \times n$  row matrix multiplied by a  $n \times 1$  column matrix:

$$(a_{11} \quad a_{12} \quad \dots \quad a_{1n}) \begin{pmatrix} b_{11} \\ b_{21} \\ \mathbf{M} \\ b_{n1} \end{pmatrix} =$$

## Revenue of a car dealer

- A car dealer sells four model types: A,B,C,D. On a given week, this dealer sold 10 cars of model A, 5 of model B, 8 of model C and 3 of model D. The selling prices of each automobile are respectively \$12,500, \$11,800, \$15,900 and \$25,300. Represent the data using matrices and use matrix multiplication to find the total revenue.

## Solution using matrix multiplication

- We represent the number of each model sold using a row matrix (4X1) and we use a 1X4 column matrix to represent the sales price of each model. When a 4X1 matrix is multiplied by a 1X4 matrix, the result is a 1X1 matrix of a single number.

$$\begin{bmatrix} 10 & 5 & 8 & 3 \end{bmatrix} \begin{bmatrix} 12,500 \\ 11,800 \\ 15,900 \\ 25,300 \end{bmatrix} = [10(12,500) + 5(11,800) + 8(15,900) + 3(25,300)] = [387,100]$$

## Matrix Product

- If  $\mathbf{A}$  is an  $m \times p$  matrix and  $\mathbf{B}$  is a  $p \times n$  matrix, the **matrix product** of  $\mathbf{A}$  and  $\mathbf{B}$  denoted by  $\mathbf{AB}$  is an  $m \times n$  matrix whose element in the  $i$ th row and  $j$ th column is the real number obtained from the product of the  $i$ th row of  $\mathbf{A}$  and the  $j$ th column of  $\mathbf{B}$ .
- If the number of columns of  $\mathbf{A}$  does **not equal** the number of rows of  $\mathbf{B}$ , the matrix product  $\mathbf{AB}$  is **not defined**.

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \\ 10 & 11 & 12 \end{pmatrix} = \begin{pmatrix} 70 & 80 & 90 \\ 158 & 184 & 210 \end{pmatrix}$$

## Undefined matrix multiplication

Why is this matrix multiplication not defined?

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \\ 10 & 11 & 12 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{pmatrix} \text{ is not defined}$$

## More examples:

$$\begin{bmatrix} 3 & 1 & -1 \\ 2 & 0 & 3 \end{bmatrix} \begin{bmatrix} 1 & 6 \\ 3 & -5 \\ -2 & 4 \end{bmatrix}$$

## Is Matrix Multiplication Commutative?


$$\begin{bmatrix} 1 & 6 \\ 3 & -5 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 3 & 1 & -1 \\ 2 & 0 & 3 \end{bmatrix}$$

## Practical application

- Suppose you are a business owner and sell clothing. The following represents the number of items sold and the cost for each item: Use matrix operations to determine the total revenue over the two days:
- Monday: 3 T-shirts at \$10 each, 4 hats at \$15 each, and 1 pair of shorts at \$20.  
Tuesday: 4 T-shirts at \$10 each, 2 hats at \$15 each, and 3 pairs of shorts at \$20.

## Solution of practical application

- Represent the information using two matrices: The product of the two matrices give the total revenue:



- Then your total revenue for the two days is  $= [110 \quad 130]$   
Price Quantity=Revenue

## Practical application

- A company with manufacturing plants located in Mass. And Virginia has labor-hour and wage requirements for the manufacture of three types of boats: What is MN?

$$\begin{array}{l}
 \text{1 man boat} \\
 \text{2 man boat} \\
 \text{3 man boat}
 \end{array}
 \begin{array}{c}
 \begin{array}{ccc}
 \text{Cutting} & \text{Assemb.} & \text{Package.} \\
 \text{Dept.} & \text{Dept.} & \text{Dept.}
 \end{array} \\
 \begin{pmatrix}
 .6 & .6 & .2 \\
 1 & .9 & .3 \\
 1.5 & 1.2 & .4
 \end{pmatrix} = M
 \end{array}
 \quad
 N = \begin{array}{c}
 \begin{array}{cc}
 \text{MA} & \text{VA}
 \end{array} \\
 \begin{pmatrix}
 17 & 14 \\
 12 & 10 \\
 10 & 9
 \end{pmatrix}
 \begin{array}{l}
 \text{Cutting} \\
 \text{Dept.} \\
 \text{Assemb.} \\
 \text{Dept.} \\
 \text{Package.} \\
 \text{Dept.}
 \end{array}
 \end{array}$$