

## MATH 250: FINAL REVIEW, FALL 2019

13.5) Lines and Planes in Space. 1-10, 11-26, 27-30, 31-37, 43-58, 61-64, 65-68, 71-72, 73-76, 77-80.

- Find the equation of the line that is perpendicular to the plane  $2x + 3y - z = 5$  and contains the point  $\langle 1, -1, 2 \rangle$ .
  - Find an equation of the line that goes through the point  $(0, 2, 3)$  and is perpendicular to the vectors  $\vec{v} = \langle 1, 0, 1 \rangle$  and  $\vec{w} = \langle 1, 2, 0 \rangle$ .
  - Find the equation of the line contained in the planes and  $x + y + z = 1$  and  $2x + 3y + z = 4$ .
- Find the equation of the plane that contains the point  $(2, 3, 1)$ , and is perpendicular to the line  $\vec{r}(t) = \langle -1 + 5t, 7 - t, 3t \rangle$ .
  - Find the equation of the plane that contains the points  $(1, 2, 0)$ ,  $(2, 3, 1)$ , and  $(3, 2, 1)$ .
  - Find the equation of the plane that contains the point  $(1, 2, 3)$  and contains the line  $\vec{r}(t) = \langle 3 - t, 2 + t, 1 + 2t \rangle$ .

13.6) Cylinders and Quadric Surfaces. 1-6, 7-12, 15-20, 21-28, 29-51, 54-58, 60.

- Sketch the surfaces.
  - $4x^2 + z^2 = 4$ .
  - $z = 4 - y^2$ .
- Sketch the  $xy$ ,  $xz$ , and  $yz$  traces. Then sketch the surface.
  - $x^2 - y^2 - z^2 = 1$ .
  - $-x^2 + y^2 + 4z^2 = 4$ .
  - $9x^2 - y^2 + 9z^2 = 0$ .

15.1) Graphs and Level Curves. 25-33, 34, 35, 36-43, 74-77.

- Sketch each function  $f(x, y)$ .
  - $f(x, y) = 6 - 2x - 3y$
  - $f(x, y) = x^2 + \frac{1}{4}y^2$
  - $f(x, y) = y^2 - x^2$
- Sketch the level curves  $f(x, y) = c$  for  $c = -1, 0, 1, 2$ .

- (a)  $f(x, y) = 2x - y$ .
- (b)  $f(x, y) = x^2 - y^2$ .
- (c)  $f(x, y) = 3 - e^{x^2+y^2}$ .

15.2) Limits and Continuity. 10-12, 13-27, 29-34, 35-50, 52-53, 62-67, 71.

1. (a) Find  $\lim_{(x,y) \rightarrow (-1,2)} \frac{\ln(x+y)}{x^2+y^2}$ .
- (b) Find  $\lim_{(x,y) \rightarrow (3,-1)} \frac{x^2-9y^2}{xy+3y^2}$ .
2. (a) Show that  $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2+2y^2}{2x^2+y^2}$  does not exist.
- (b) Show that  $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2y}{x^3+xy^2}$  does not exist.

15.3) Partial Derivatives. 1-9, 11-14, 15-30, 32-34, 38-46, 48-53, 54-59.

1. Use the **limit definition** to find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$ .
  - (a)  $f(x, y) = 3x - y$ .
  - (b)  $f(x, y) = xy^2$ .
2. Find all first and second partials.
  - (a)  $f(x, y) = \sin(xy)$ .
  - (b)  $f(x, y) = \ln(x^2 + y^3)$ .

15.4) The Chain Rule. 9-18, 19-26, 27-28, 29-30, 35-40, 57-59, 65, 67-69, 72-73, 75.

1.  $w = x^3y^2$ ,  $x = t^2 + 1$ ,  $y = t - e^{2-t}$ . Use the **chain rule** to find  $\frac{dw}{dt}$  at  $t = 2$ .
2.  $w = \ln(xy)$ ,  $x = 2u + 3v$ ,  $y = \frac{v^2}{u}$ . Use the **chain rule** to find  $\frac{\partial w}{\partial u}$  and  $\frac{\partial w}{\partial v}$  at  $(u, v) = (2, -1)$ .
3. (a) If  $F(x, y, z) = c$  where  $c$  is constant and  $y = y(x, z)$ , use the chain rule to show that  $\frac{\partial y}{\partial z} = -\frac{\partial F}{\partial z} / \frac{\partial F}{\partial y}$ .
- (b) Use this formula to find  $\frac{\partial y}{\partial z}$ , when  $ye^{xz} + xe^{yz} = 1$ .
4. Find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  by implicit differentiation.  $x^2y + y^2z + xz^3 = 1$ .
5. For a unit-length pendulum, if  $\theta$  is the angular position and  $v = \frac{d\theta}{dt}$ , then  $\frac{dv}{dt} = -g \sin(\theta)$ , where  $g$  is constant. Use the chain rule to show that  $\frac{dE}{dt} = 0$ , where  $E(\theta, v) = \frac{1}{2}v^2 - g \cos(\theta)$ .

15.5) Directional Derivatives and Gradient. 1-10, 11-12. 13-20, 21-30, 31-36, 43-44, 47-50, 59-64, 69-72, 74, 75-78, 81, 82, 85.

1. Find the gradient of  $f$  at the point  $P$ . Then find  $D_{\vec{u}}f$  in the direction  $\vec{u}$ .  $f(x, y, z) = x^2y + z^3$ ,  $P = (-1, 2, 1)$ ,  $\vec{u} = \langle \frac{-2}{3}, \frac{2}{3}, \frac{1}{3} \rangle$ .
2.  $f(x, y) = x^2 + xy - y^3$ .
  - (a) Find the maximum rate of change of  $f$  (steepest ascent), and the direction of the maximum rate of change, at  $P = (3, 2)$ .
  - (b) Find a vector that points in a direction of no change of  $f$ , at  $P = (3, 2)$ .
3.  $f(x, y) = xy$ .
  - (a) Graph the level set of  $f$  through the point  $(2, 1)$ .
  - (b) Include the vector  $\nabla f(2, 1)$  on your graph.
  - (c) Find the tangent line to the level set at the point  $(2, 1)$  and include it on your graph.
  - (d) How are the answers to parts b and c related?

15.6) Tangent Planes. 3-4, 9, 11, 13-28, 29-32, 54-56. |

1. Find the tangent plane to the surface at the given point.
  - (a)  $3x^2 + xy + z^2 = 5$ ,  $P = (-1, 2, 2)$ .
  - (b)
  - (c)  $xe^{yz} = 3$ ,  $P = (3, 0, 2)$ .
  - (d)  $\frac{x-y}{3y+z} = 1$ ,  $P = (3, 1, -1)$ .
2. Find the tangent plane to the surface at the given point.
  - (a)  $z = \sqrt{x^2 - y^2}$ ,  $P = (5, 4, 3)$ .
  - (b)  $z = 3 - \sin(xy)$ ,  $P = (2, 0, 3)$ .
  - (c)  $y = xe^{x+2z}$ ,  $P = (2, 1, -1)$ .

15.7) Maximum/Minimum Problems. 9-12, 13-22, 23-37, 41-42, 43-46, 62-66, 71. |

1. Find any critical points and classify each as relative maximum, relative minimum, or saddle.
  - (a)  $f(x, y) = e^{-x^2} + e^{-y^2}$
  - (b)  $f(x, y) = x^2 + y^2 + xy - x - 2y$

- (c)  $f(x, y) = xy - 2x - y$   
 (d)  $f(x, y) = x^4 - 2x^2 + y^2$

- Find the minimum of  $x^2 + y^2 + z^2$  if  $(x, y, z)$  is on the plane  $x - z = 2$ . Use the second derivative test to prove your answer is a local minimum.
- Find the maximum volume of a box  $V = xyz$  if the point  $(x, y, z)$  is on the paraboloid  $z = 4 - x^2 - y^2$ . ( $x, y, z > 0$ .) Use the second derivative test to prove your answer is a local maximum.

15.8) Lagrange Multipliers. 3-4, 5-6, 7-23, 26, 27-36. |

- Use Lagrange multipliers to find the maximum and minimum of  $f(x, y) = xy^3$  if  $x^2 + y^2 = 4$ .
- The area of a rectangle with vertices  $(\pm x, \pm y)$  is  $4xy$ . Use Lagrange multipliers to find the maximum area of such a rectangle with vertices on the ellipse  $4x^2 + y^2 = 32$ .
- Use Lagrange multipliers to find the minimum distance between the plane  $3x + y - z = 18$  and the point  $(2, 0, -1)$ . (Hint: To find  $(x, y, z)$  you can minimize the square of the distance,  $f(x, y, z) = (x - 2)^2 + y^2 + (z + 1)^2$ .)

16.1) Double Integrals, Rectangular Regions. 1-3, 5-6, 7-24, 25-35, 36-39, 40-45, 46-50, 53-54. |

- Find the average value of  $f(x) = 2y - x$  for  $0 \leq x \leq 2$ ,  $1 \leq y \leq 3$ .
- Choose the most convenient order of integration and evaluate the integral.
  - $\iint_R xye^{xy^2} dA$ ,  $0 \leq x \leq \ln(3)$ ,  $0 \leq y \leq 1$ .
  - $\iint_R \frac{y}{\sqrt{1+xy}} dA$ ,  $0 \leq x \leq 1$ ,  $0 \leq y \leq 3$ .

16.2) Double Integrals, General Regions. 5-8, 9-10, 11-27, 28-34, 35-42, 43-53, 57-62, 63-68, 69, 70, 71, 73-80, 85-90, 95-96, 99-102.

- Find the volume under  $f(x, y) = 2y - 1$ , on the region between  $x = y^2 + 3$  and  $x = 3y + 1$ .
- Evaluate the integrals by changing the order of integration.
  - $\int_0^6 \int_{\frac{1}{2}x}^3 e^{y^2} dy dx$ .
  - $\int_0^4 \int_0^{\sqrt{y}} e^{12x-x^3} dx dy$

16.3) Double Integrals in Polar Coordinates. 7-10, 11-14, 15-18, 19-20, 21-30, 31-40, 42, 44-46, 47, 49-50, 53-54, 57-60, 65-68.

1. Evaluate the integral by converting to polar.

(a)  $\int_0^4 \int_0^{\sqrt{16-x^2}} xy \, dydx.$

(b)  $\int_{-2}^2 \int_0^{\sqrt{4-y^2}} \frac{1}{(1+x^2+y^2)^2} \, dx dy.$

(c)  $\int_1^2 \int_{-x}^x 1 \, dydx.$

16.4) Triple Integrals. 4-6, 7-14, 15-29, 30-35, 36-37, 38-46, 47-50, 51-54, 57-58, 62-63, 67-70.

- Express the volume under  $x^2 + 4y + z = 4$ , with  $x, y, z \geq 0$  by six different triple integrals.
- $V$  is the region between  $y = x^2$ ,  $z = 0$ , and  $y + 2z = 4$ .
  - Find the volume using an integral  $dydx dz$ .
  - Find the volume using an integral  $dzdxdy$ .

16.5) Cylindrical and Spherical Coordinates. 3-4, 9-10, 11-14, 15-22, 23-28, 29-34, 35-38, 41-47, 48-54, 58-61, 62-63, 64-65, 66-72, 77-79.

- Evaluate  $\iiint_V \frac{z}{(x^2+y^2)^{\frac{3}{2}}} \, dx dy dz$  where  $V$  is the region with  $1 \leq x^2 + y^2 \leq 4$  and  $0 \leq z \leq 4 - x^2 - y^2$ .
- Evaluate  $\iiint_V xz \, dx dy dz$  where  $V$  is the region inside the sphere of radius 2 in the first octant.
- Find the volume of the region above the cone  $z = \sqrt{x^2 + y^2}$  and below  $z = 3$  by an integral in cylindrical coordinates.
  - Find the same volume using an integral in spherical coordinates.
- Find the volume of the region inside the sphere  $x^2 + y^2 + z^2 = 25$  and outside the cylinder  $x^2 + y^2 = 9$  by an integral in cylindrical coordinates.
  - Find the same volume using an integral in spherical coordinates.

16.7) Change of Variables. 5-11, 13-16, 17-22, 23-26, 27-30, 31-36, 37-39, 41-44, 46-47, 48, 50-52, 53, 56.

- Let  $R$  be the region between  $xy = 1$ ,  $xy = 2$ ,  $y = x$ ,  $y = 3x$ . Use the change of variables  $u = xy$ ,  $v = \frac{y}{x}$  to find the area of  $R$  by a double integral.
- Let  $R$  be the region with  $1 \leq x + 2y \leq 3$ ,  $0 \leq 3x + 4y \leq 2$ . Use a change of variables to evaluate  $\iint_R x \, dA$ .

3. Let  $R$  be the region inside the ellipse  $\frac{x^2}{4} + \frac{y^2}{25} = 1$ , with  $y \geq 0$ . Evaluate  $\int_R y \, dx \, dy$  by making the change of variables  $x = 2u, y = 5v$ .

17.1) Vector Fields. 2, 8-15, 18, 24, 25-30, 35-42, 43-45, 47-48, 49-52. |

1. Sketch the vector fields:  $\vec{F}(x, y) = \langle x, y \rangle$  and  $V(x, y) = \langle y, -x \rangle$ .
2. Plot the vector field at the points  $(1, 0)$ ,  $(0, 1)$ ,  $(1, 1)$ , and  $(-1, 1)$ .  $\vec{F}(x, y) = \langle 2x + y, -x + 2y \rangle$ .

17.2) Line Integrals. 4-10, 12-16, 17-34, 35-36, 39-40, 41-46, 47-48, 49-56, 57-60, 62, 64-65, 68, 70-72, 73.

1. Evaluate the line integrals.
  - (a)  $\int_c yz \, dx + xyz \, dz$   $\vec{r}(t) = \langle 1, t, t^2 \rangle$ , from  $(1, 0, 0)$  to  $(1, 2, 4)$ .
  - (b)  $\int_c xz \, ds$ , where  $c$  is the line from  $(0, 1, -1)$  to  $(2, 0, 1)$ .
  - (c)  $\int_c \langle x - y, 2y \rangle \cdot d\vec{r}$ , where  $c$  is the curve along  $x = y^4$  that connects  $(1, -1)$  to  $(16, 2)$ .

17.3) Conservative Vector Fields. 7, 9-16, 17-30, 31-34, 39-42, 44, 45-50, 51-52, 54-56, 59-62, 63-64. |

1.  $\vec{F}(x, y) = \left\langle \frac{x}{\sqrt{x^2+y^2}}, \frac{y}{\sqrt{x^2+y^2}} \right\rangle$ .
  - (a) Find a function  $\phi$  such that  $\nabla\phi = \vec{F}$ .
  - (b) Use this to find  $\int_c \vec{F} \cdot d\vec{r}$ , where  $\vec{r}(t) = \langle 1 + t, 2t \rangle$ ,  $0 \leq t \leq 2$ .
2.  $\vec{F} = \langle 1 + \cos(y), 2y - x \sin(y) \rangle$ .
  - (a) Show  $\vec{F}$  is conservative.
  - (b) Find a function  $\phi$  such that  $\nabla\phi = \vec{F}$ .
  - (c) Use this to evaluate  $\int_c \vec{F} \cdot d\vec{r}$ ,  $\vec{r}(t) = \langle 2^t, \pi t \rangle$ ,  $0 \leq t \leq 1$ .
3. Show  $\vec{F}$  is conservative, and find a function  $\phi$  such that  $\nabla f = \vec{F}$ .  
 $\vec{F} = \langle 2x + y, x + 2, 3z^2 \rangle$ .

17.4) Green's Theorem. 9-14, 15-16, 17-20, 21-25, 27-30, 31-40, 41-45, 48, 53-54. |

1. Use Green's theorem to compute  $\int_c \vec{F} \cdot d\vec{r}$  where  $\vec{F} = (x^2 + y^2)\vec{i} + 2xy\vec{j}$ , and  $c$  along the rectangle with vertices  $(0, 0)$ ,  $(2, 0)$ ,  $(2, 1)$ ,  $(0, 1)$  oriented counterclockwise.

2. Let  $R$  be the region  $x^2 \leq y \leq 1$ ,  $\vec{F} = \langle x - y, x + y \rangle$ .
  - (a) Compute  $\int_c \vec{F} \cdot d\vec{r}$  directly, where  $c$  is the boundary oriented counterclockwise. (The boundary has two components!)
  - (b) Use Green's theorem to compute  $\int_c \vec{F} \cdot d\vec{r}$ . Show that you get the same answer.
3. Use Green's theorem to compute  $\int_c \langle x - y, 2x + 3y \rangle \cdot \vec{n} ds$ , where  $c$  is the triangle with vertices  $(0, 0)$ ,  $(1, 0)$ ,  $(1, 2)$ .
4. Verify Green's Theorem (flux form) by computing both sides.  $\vec{F} = \langle x^3, y^3 \rangle$  with  $c$  the circle which bounds the region  $x^2 + y^2 \leq 4$ , oriented counterclockwise.

17.5) Divergence and Curl. 9-16, 17, 19, 21-22, 27-34, 41, 42, 44, 65, 67-69, 73. |

1. Find  $\nabla \cdot \vec{F}$  and  $\nabla \times \vec{F}$ .  $\vec{F}(x, y, z) = \langle x^3, x^2y, yz^3 \rangle$ .
2. Show that if  $\vec{F} = \nabla\phi$  for some  $\phi(x, y, z)$ , then  $\nabla \times \vec{F} = \vec{0}$ .
3. Show there is no function  $\phi(x, y, z)$  with  $\nabla\phi = \langle x^2 + y^2, xyz, e^{xz} \rangle$ .

17.6) Surface Integrals. 9-14, 15, 17-18, 19-24, 25-28, 29-34\*, 35-38\*, 43-48\*, 52, 70-72, 74-75. |

\*: For 29-34, 35-38, 43-48, you **must** parametrize the surface by  $r(u, v)$  and use the parametric form to do the integrals, rather than the 'explicit' form indicated in the book here.

1. Evaluate  $\iint_S 1 dS$ , where  $S$  is the paraboloid  $z = 1 - x^2 - y^2$  with  $z \geq 0$ .
2. Evaluate  $\iint_S \langle 1, 0, 2 \rangle \cdot d\vec{S}$ , where  $S$  is the cone  $z = \sqrt{x^2 + y^2}$  with  $0 < z < 2$ . Upward pointing normal.
3. Use a surface integral to find the area of the region of the plane  $z = x + 2y + 3$  with  $x^2 \leq y \leq 3x$ .
4. A surface of revolution given by  $y = f(x)$  revolved around the  $x$ -axis can be written  $\vec{r}(u, v) = \langle v, f(v) \cos(u), f(v) \sin(u) \rangle$ , with  $0 \leq u \leq 2\pi$ ,  $a \leq v \leq b$ . Use this to derive the formula  $\text{Area} = 2\pi \int_a^b f(v) \sqrt{1 + (f'(v))^2} dv$ .
5. A helicoid is given by the parametric surface  $\vec{r}(u, v) = \langle u \cos(v), u \sin(v), v \rangle$ , with  $0 \leq u \leq 1$ ,  $0 \leq v \leq \pi$ . Evaluate the integrals.
  - (a)  $\iint_S \langle x, y, z \rangle \cdot d\vec{S}$ . Upward pointing normal.
  - (b)  $\iint_S y dS$ .

17.7) Stokes' Theorem. 5-10, 11-16, 17-24, 30-33, 45. |

1. Compute both sides in Stokes' theorem and show that they are equal, for the surface  $z = 4 - x^2 - y^2$ ,  $z \geq 0$ .  $\vec{F} = \langle -y, x, z^2 \rangle$ .
2. Use Stokes' theorem to evaluate  $\iint_S \nabla \times \vec{F} \cdot d\vec{S}$ , where  $S$  is the hemisphere  $x^2 + y^2 + z^2 = 9$ ,  $y \geq 0$ , and  $\vec{F} = \langle x - z, e^{xy}, x + z \rangle$ . Right pointing normal.
3. Use Stokes' theorem to evaluate  $\int_c \vec{F} \cdot d\vec{r}$  where  $c$  is the triangle with vertices  $(1, 0, 0)$ ,  $(0, 1, 0)$ ,  $(0, 0, 1)$  oriented counterclockwise, and  $\vec{F} = \langle x - y, x + y, z \rangle$ . (Hint:  $z = 1 - x - y$  gives the surface, with  $0 \leq x \leq 1$ ,  $0 \leq y \leq 1 - x$ ).

17.8) Divergence Theorem. 9-12, 13-16, 17-24, 25-27, 30. |

1.  $\vec{F}(x, y, z) = \langle y, -x, z^2 \rangle$ . Evaluate both sides of the divergence theorem and show that they are equal, for the region  $x^2 + y^2 \leq z \leq 4$ . (The boundary has two pieces,  $z = x^2 + y^2$  and  $z = 4$ . Outward pointing normals.)
2. Use the divergence theorem to evaluate  $\iiint_S \vec{F} \cdot d\vec{S}$ . The surfaces have outward pointing normal.
  - (a)  $\vec{F} = \langle x, y, z \rangle$ ,  $S$  the boundary of the tetrahedron  $x, y, z \geq 0$ ,  $x + 2y + 3z \leq 6$ .
  - (b)  $\vec{F} = \langle x^3z, y^3z, xy \rangle$ .  $S$  the sphere  $x^2 + y^2 + z^2 = 9$ .